


Pinpoint®

F U N G I C I D E



TURFGRASS FUNGICIDE
A SYSTEMIC FUNGICIDE FOR CONTROL
OF CERTAIN TURFGRASS DISEASES

Active Ingredient	By Wt
Mandestrobin*	43.4%
Other Ingredients	56.6%
Total	100.0%

* (RS)-2-methoxy-N-methyl-2-[α -(2,5-xylyloxy)-*o*-tolyl] acetamide

EPA Reg. No. 59639-199 EPA Est. 5481-ID-1[®], 228-IL-2[®]
 Superscript is first letter of lot number.

Pinpoint® Fungicide is a suspension concentrate containing 4 lb active ingredient per gallon.

NET CONTENTS 60 FLUID OUNCES

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE NEXT PAGE FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

FIRST AID

If swallowed: Call a poison control center or doctor immediately for treatment advice.

Have person sip a glass of water if able to swallow.

Do not induce vomiting unless told to by a poison control center or doctor.

Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 800-892-0099 for emergency medical treatment information.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, socks and shoes.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic organisms. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposing of equipment washwaters or rinsate. Drift and runoff from treated areas may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is coveralls, chemical-resistant gloves made of any waterproof material, socks and shoes.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to the uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, greenhouses or sod farms.

Keep all unprotected persons out of operating areas or vicinity where there may be drift.

Do not enter or allow others to enter treated areas until sprays have dried.

**DISCLAIMER, RISKS OF USING THIS PRODUCT,
LIMITED WARRANTY
AND LIMITATION OF LIABILITY**

IMPORTANT: Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty, and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as “Buyer”) of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application, (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, **THEN THIS PRODUCT SHOULD NOT BE APPLIED.** By applying this product Buyer acknowledges and accepts these inherent unintended risks **AND AGREES THAT TO THE**
(continued)

(continued)

EXTENT CONSISTENT WITH APPLICABLE LAW, RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

To the extent consistent with applicable law, Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. To the extent consistent with applicable law, Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

LIMITED WARRANTY

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label **and subject to the Risks of Using This Product as described above. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, EXCEPT AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED.** No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

To the extent consistent with applicable law, Valent or Seller shall not be liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE BUYER, AND THE

(continued)

(continued)

EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PROMPT NOTICE OF CLAIM

Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is later, so that an immediate inspection of the affected property and growing crops can be made.

To the extent consistent with applicable law, if Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing Disclaimer, Risks of Using This Product, Limited Warranty and Limitation of Liability, which may not be modified by any oral or written agreement.

TANK MIX

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor, consistent with applicable law.

Read and follow the entire label of each product to be used in the tank mix with this product.

TABLE OF CONTENTS

Product Information	8-12
Mode of Action	9
Restrictions	9
Precaution	9
Resistance Management	9-10
Rainfastness	10
Tank Mixtures	10
Jar Test to Determine Compatibility of Adjuvants and <i>Pinpoint</i> Fungicide. ...	11
Sprayer Preparation	11
Application Equipment	11
Sprayer Cleanup	12
Mixing Instructions	12
Turfgrass	12-16
Listed Turfgrass Species	12
Application Directions	14
Turfgrass Tolerance	14
Restrictions	14
Turfgrass - Specific Diseases, Rates, Timing and Application Information table	15-16
Storage and Disposal	17

PRODUCT INFORMATION

Pinpoint Fungicide is a broad spectrum fungicide for control of certain diseases in turfgrass. *Pinpoint* Fungicide has preventive, curative and systemic properties. *Pinpoint* Fungicide can be applied pre- or post-infection, but is most effective when applied prior to infection. Optimal disease control is achieved when *Pinpoint* Fungicide is applied in a regularly scheduled spray program used in combination and/or rotation with other effective fungicides that have different modes of action and are effective against the same target pathogen(s).

MODE OF ACTION

Pinpoint Fungicide belongs to the strobilurin or Quinone Outside Inhibitors (QOI) class (Group 11) of fungicides as classified by the Fungicide Resistance Action Committee (FRAC).

RESTRICTIONS

- Not for Sale, Sale Into, Distribution and/or Use in Nassau and Suffolk Counties of New York State.
- Do not apply this product within 50 feet of any water bodies that connect with any off-site water bodies such as a freshwater lake, pond, river, stream or wetland in the State of New York.

PRECAUTION

Always mix product thoroughly before use.

Resistance Management

For resistance management, *Pinpoint* Fungicide contains a Group 11 fungicide. Any fungal population may contain individuals naturally resistant to *Pinpoint* and other Group 11 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of *Pinpoint* Fungicide or other Group 11 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.

(continued)

(continued)

- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Valent U.S.A. LLC at 800-89-Valent (898-2536). You can also contact your pesticide distributor or university extension specialist to report resistance.

RAINFASTNESS

Pinpoint Fungicide is rainfast 2 hours after application. Disease control may be reduced if rainfall occurs within 2 hours after application.

TANK MIXTURES

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

JAR TEST TO DETERMINE COMPATIBILITY OF ADJUVANTS AND *PINPOINT* FUNGICIDE

Perform a jar test before mixing commercial quantities of *Pinpoint* Fungicide, when using this product for the first time, when using new adjuvants, when using new tank mixes, or when using a new water source.

1. Add 1 pt of the water to a quart jar. Use water from the same source and temperature as that to be used in the spray tank mixing operation.
2. Add 2 g of *Pinpoint* Fungicide to the quart jar, gently mix until product goes into suspension.
3. Add 1 ml of new adjuvant and/or appropriate amount of new tank mix partner and gently mix.
4. Place cap on jar, invert 10 times, let stand for 15 minutes, evaluate.
5. An acceptable tank mix combination will have a smooth, uniform appearance. If any of the following conditions are observed, question the choice of spray mix components:
 - a) Layer of oil or globules on the mixture's surface.
 - b) Flocculation: formation of fluffy, cloudlike aggregates or masses in suspension or as a layer on the bottom of the jar.
 - c) Clabbering: thickening texture (coagulated) like gelatin or cottage cheese.

SPRAYER PREPARATION

Before applying *Pinpoint* Fungicide, start with clean, well maintained application equipment. The spray tank hoses and booms must be cleaned to ensure no residue from previous spraying operations remains in the sprayer. The spray equipment must be cleaned according to the manufacturer's directions for the last product used before the equipment is used to apply *Pinpoint* Fungicide. If two or more products were tank mixed prior to *Pinpoint* Fungicide application, the most restrictive cleanup procedure must be followed.

APPLICATION EQUIPMENT

Application equipment must be clean and in good repair. Check nozzles frequently for accuracy.

SPRAYER CLEANUP

Clean sprayer equipment each day following *Pinpoint* Fungicide application. After application is complete, use the following steps to clean spray equipment:

1. Completely drain the spray tank, rinse the sprayer thoroughly, including the inside and outside of the tank and all in-line screens.
2. Fill the spray tank with clean water and flush all hoses, booms, screens and nozzles.
3. Drain tank completely.
4. Remove all nozzles and screens and rinse them in clean water.

MIXING INSTRUCTIONS

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. While agitating, slowly add the *Pinpoint* Fungicide to the spray tank. Agitate to create a rippling or rolling action on the water surface.
3. If tank mixing *Pinpoint* Fungicide with other labeled pesticides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates and then solutions.
4. Add any required adjuvants.
5. Fill spray tank to desired level with water. Continue agitation until all spray solution has been applied.
6. Mix only the amount of spray solution that can be applied the day of mixing. Apply *Pinpoint* Fungicide within 24 hours of mixing.

TURFGRASS

LISTED TURFGRASS SPECIES

Annual Bluegrass
Bentgrass
Bermudagrass
Centipedegrass

Fescue
Kentucky Bluegrass
Perennial Ryegrass
Zoysia

USE SITES

Pinpoint Fungicide may be used for disease control on turfgrass in golf courses, lawns and turfgrass in landscape areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas, athletic fields and sod farms.

SPECIAL INSTRUCTIONS

- Use *Pinpoint* Fungicide in conjunction with turf management practices that promote good plant health and optimum disease control. Proper identification of disease infestations is essential for providing control. Consult diagnostic guides, kits or other means of identification to determine the best control measures.
- *Pinpoint* Fungicide is more effective when used preventively or early in disease development. Length of disease control will vary according to local environmental conditions, disease pressure and turfgrass management practices.
- Apply to dry turf or after mowing. For control of foliar diseases, allow turf to dry before irrigation. For control of soil borne diseases, *Pinpoint* Fungicide can be watered in immediately after application or applied with a high volume of water.
- *Pinpoint* Fungicide may be tank mixed with other registered turfgrass fungicides. Perform a jar test as described above to determine compatibility before making applications. Directions for use of *Pinpoint* Fungicide have been developed using no adjuvant in the spray solution.
- Important: Under conditions of optimal disease development (such as high humidity and temperature), use higher labeled rates and shorter application intervals. When applications are made curatively, use higher rates of *Pinpoint* Fungicide and shorter application intervals. Use higher application rate in areas with a history of severe disease pressure.
- Failure to follow label directions may result in turfgrass injury and/or inferior disease control.

APPLICATION DIRECTIONS

Apply *Pinpoint* Fungicide using standard, low-pressure spray equipment designed to deliver coarse to medium size spray droplets (do not use nozzles designed to deliver fine droplets). Apply in a sufficient volume of water to provide thorough spray coverage and a uniform spray pattern. Do not position the spray boom more than 20 inches above the turf. For foliar disease, apply 2 gal of spray solution per 1,000 sq ft. For crown and root diseases, apply 2 to 4 gal of spray solution per 1,000 sq ft. For spot treatment, mix 0.1 to 0.2 fl oz of *Pinpoint* Fungicide per gallon of water.

TURFGRASS TOLERANCE

Due to the large number of turfgrass types, possible tank mixes and application techniques, it has not been determined if *Pinpoint* Fungicide can be safely used on all turfgrasses under all conditions. Before making a large scale application, determine safety in a small test area. Monitor the test area for at least 2 weeks to determine safety under local growing conditions.

RESTRICTIONS

- Do not apply more than 13.5 fl oz (0.42 lb ai) of *Pinpoint* Fungicide per acre per application.
- Do not make more than 4 applications of *Pinpoint* Fungicide per acre per year.
- Do not apply more than 54 fl oz (1.68 lb ai) of *Pinpoint* Fungicide per acre per year.
- Do not apply through any type of irrigation system.
- Do not apply by aerial application.
- Do not graze animals on treated areas for one year after application.
- Do not feed clippings from treated areas to livestock or poultry.

TURFGRASS – Specific Diseases, Rates, Timings and Application Information

Diseases (<i>Pathogens</i>)	Application Rates		Application Interval/ Timing	Special Instructions
	fl oz/ 1,000 sq ft	fl oz/A		
Dollar Spot (<i>Sclerotinia homoeocarpa</i>)	0.275 to 0.31	12 to 13.5	14 to 21 days	Use preventive-ly. Begin application when conditions favor disease development. Do not make more than 2 <i>Pinpoint</i> Fungicide applications without alternating to a non-Group 11 fungicide for resistance management.
Brown Patch (<i>Rhizoctonia solani</i>) (suppression only) Fairy Ring (various <i>Basidiomycetes</i>) Rust Diseases (<i>Puccinia</i> spp.) Take-all Patch (<i>Gaeumannomyces graminis</i>)	0.31	13.5	14 days	Begin applications when conditions favor disease development or when the disease first appears. Do not make more than 2 <i>Pinpoint</i> Fungicide applications without alternating to a non-Group 11 fungicide for resistance management.

fl oz Product/ 1,000 sq ft	fl oz Product/Acre	lb ai/Acre	Maximum Applications per Year
0.275	12.0	0.375	4
0.31	13.5	0.42	4

STORAGE AND DISPOSAL

PESTICIDE STORAGE

Store in a cool dry place. Keep pesticide in original container. Keep container closed when not in use. Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Do not put dilute into food or drink containers. Do not store in or around the home.

For help with any spill, leak, fire or exposure involving this material, call day or night 800-892-0099.

PESTICIDE DISPOSAL

Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available, or dispose in a sanitary landfill or incinerate.

SPECIMEN

SPECIMEN

SPECIMEN

Pinpoint and Products That Work, From People Who Care are registered trademarks of Valent U.S.A. LLC

©2019 Valent U.S.A. LLC

Information contained in this booklet is accurate at the time of printing. Since product testing is a continuous process, please read and follow the directions on the product label for the most current directions and precautionary statements.

Always check with your state to verify state registration status or call 800-89-VALENT (898-2536).

Manufactured for:

Valent U.S.A. LLC

P.O. Box 8025

Walnut Creek CA 94596-8025

Form 2069-F

Made in U.S.A.

EPA Reg. No. 59639-199

EPA Est. 5481-ID-1[®], 228-IL-2[®]

Superscript is first letter of lot number.

059639-00199.20190821.S-2200_4SC_VPP.AMEND.NY.FPL

SAL20190826

Distributed by:



Nufarm

Grow a better tomorrow.

Nufarm Americas Inc.

11901 S. Austin Ave.

Alsip, IL 60803



For state registration and/or supplemental labels, please call or visit us online.

Products That Work, From People Who Care[®]

valentpro.com | 800-89-VALENT (898-2536)

Always read and follow label instructions.


©2019 Valent U.S.A. LLC. All rights reserved.

Printed in the U.S.A.

MANDESTROBIN GROUP 11 FUNGICIDE

Pinpoint®

F U N G I C I D E



TURFGRASS FUNGICIDE
A SYSTEMIC FUNGICIDE FOR CONTROL
OF CERTAIN TURFGRASS DISEASES

Active Ingredient	By Wt
Mandestrobin*	43.4%
Other Ingredients	56.6%
Total	100.0%

* (RS)-2-methoxy-N-methyl-2-[α -(2,5-xylyloxy)-o-tolyl] acetamide

EPA Reg. No. 59639-199 EPA Est. 5481-ID-1[®], 228-IL-2[®]
Superscript is first letter of lot number.

Pinpoint® Fungicide is a suspension concentrate containing 4 lb active ingredient per gallon.

**KEEP OUT OF
REACH OF CHILDREN
CAUTION**
**SEE BOOKLET FOR ADDITIONAL
PRECAUTIONARY STATEMENTS.**

NET CONTENTS 60 FLUID OUNCES

Form 2069-F
Date 11.19.19